

Volume 3, Issue 2, 2014

October 2014

ISSN: 2319-880X

# International Journal Of Agricultural Science & Technology




This Journal is an academic and peer-reviewed publication (Print ISSN 2319-880X )

An International Refereed Journal

**Academic &  
RESEARCH  
Publications**

ARP. Give your IDEAS an Expression

[www.manishanpp.com](http://www.manishanpp.com)





## ABOUT THE JOURNAL

**INTERNATIONAL JOURNAL OF AGRICULTURAL SCIENCE AND TECHNOLOGY** is a biannual academic and peer-reviewed Journal published by ACADEMIC AND RESEARCH PUBLICATIONS. It was published from year i.e. 2012. The ISSN of the JOURNAL is 2319-880X.

## OBJECTIVE OF THE JOURNAL

To promote and encourage specially YOUNG SCIENTISTS to take active part in research and get acquainted with the latest development and research in the field of environmental science and technology. To promote cooperation in the pursuit of research in general and to exchange and contribute to the progress in the field of Agricultural Science and technology in particular.

## CALL FOR PAPERS

We invite you to submit high quality papers for review and possible publication in all areas of Agricultural Science and technology which includes Soil Science, Agri. Entomology, and Gnd.water Pollution, Limnology, Toxicology, Animal Husbandry, Biological and Chemical Control, Seed Science and Technology, Agricultural Waste, Any other related topics. All authors must agree on the content of the manuscript and its submission for publication in this Journal before it is submitted to us. Manuscripts should be submitted by e-mail to the Editor at [manisha\\_npp@yahoo.com](mailto:manisha_npp@yahoo.com).

## COVERAGE OF THE JOURNAL

- |                                     |                               |
|-------------------------------------|-------------------------------|
| 1. Soil Science                     | 2. Water & Soil Pollution     |
| 3. Agricultural Entomology          | 4. Aquaculture                |
| 5. Groundwater Pollution            | 6. Pesticides                 |
| 7. Limnology                        | 8. Sericultural Biotechnology |
| 9. Toxicology                       | 10. Plant Protection Science  |
| 11 Animal Husbandry.                | 12. Meteorology               |
| 13. Biological and Chemical Control | 14. Agronomical Science       |
| 15. Seed Science and Technology     | 16. Forest and Forestry       |
| 17. Agricultural Waste              | 18. Any other related topics  |

## TYPES OF PAPERS ARE INVITED

Following types of papers are invited for publication in this Journal :-

- |  |                  |
|--|------------------|
| a) Original Research Papers of Scientific values | b) Review Papers |
| c) Short Communications                          | d) Case Reports  |
| e) Letters to the Editor                         | f) As you see    |

## REVIEWERS PROCESS

All manuscripts are reviewed by an editor and members of the Editorial Board or qualified outside REVIEWERS. Decisions will be made as rapidly as possible and the Journal strive to return reviewer's comments to authors within 6 weeks. The Editorial Board will re-review manuscripts that are accepted pending reviewers. It is the goal of the this Journal to publish manuscripts within 4 weeks after submission after getting O K report from the Author.

## CONTACT US

For quick reply, please note change of address and contact them directly by Post or email:-

For publication of your article, Acceptance letter, Review Reports, Status Report, and all other queries related to your articles, should be sent directly to the Editor-in-Chief, whose address is as follows:

**Prof. Manik Sinha, THE EDITOR-IN-CHIEF**

Email : [manik.sinha2@gmail.com](mailto:manik.sinha2@gmail.com), Contact at : 09415155631

For all publication matters related to the Journals Acceptance letter for publication of articles, Invoice, Reprints etc. should be sent directly to the PUBLICATION EDITOR whose address is as follows :

To,

*Ex. Manisha Verma, B. Sc., B. Tech.*

**Publication Editor (Chief Executive Director)**

**Academic And Research Publications**

H. Office: EC 41, Maya Enclave, New Delhi -110064

Email : [manisha\\_npp@yahoo.com](mailto:manisha_npp@yahoo.com) [www.manishanpp.com](http://www.manishanpp.com)

# IJAST

....

## International Journal Of Agricultural Science & Technology

Volume 3, Issue 2

October 2014

Cite this volume as 3(2)IJAST(2014) and so on....

“ This journal is Indexed/abstracted in Indian Science Abstract ”

© Journal on Agricultural Science and Technology. All rights reserved. No portion of material can be reproduced in part or full without the prior permission of the Editor.

Note : The views expressed herein are the opinions of contributors and do not reflect the stated policies of the

**ACADEMIC AND RESEARCH PUBLICATIONS**


**This Journal is an academic and peer-reviewed publication (Print ISSN 2319-880X )**

An International Refereed Journal



**ARP. Give your IJAST an Expression**

[www.manishanpp.com](http://www.manishanpp.com)



**Academic And Research Publications**

H. Office: EC 41, Maya Enclave, New Delhi -110064

# International Journal Of Agricultural Science & Technology

Volume 3, 2014

October 2014

Issue 2, 2014

## Editorial Board

### Editor-in-Chief

**Prof. Manik Sinha**

Former Dean, Faculty of Law,  
Dr R.M.L Awadh University, Faizabad (UP),  
Senior Advocate, Govt Of India, High Court, Lucknow  
Email: manik.sinha2@gmail.com

### Chief Editor

**Prof. Padmakar Tripathy,**  
Dept of Meteorology,  
N D University of Agri. and Tech.,  
FAIZABAD - 224 229 U P  
E-mail: pmtmausam@gmail.com

### Associate Editor

**Dr. K. K. Singh,**

Head, Agromet Services &  
Scientist-F  
India Meteorological Department  
Lodhi Road, New Delhi - 110003

**Dr. A. P. Dubey, Professor**  
(Agromet)

Division of Agrometeorology  
C.S.A. U of A & , Kanpur-208002  
Dr. S. R. Mishra Asstt. Professor  
Deptt. of Agril. Met.  
N.D.U.A. & T., Faizabad-224229  
e-mail: srmmausam@india.com

**Dr. A. K. Singh Asstt. Professor**  
Department of Crop Physiology  
N. D. U A. & T,  
FAIZABAD - 224 229  
E: assinghkumar3@gmail.com

### Associate Editors

**Mrs. Chandra Kanti Sinha**  
Retd. Laclurer in Botany(Principal Grd.),  
Govnment Collage Faizabad U.P.  
Email: chandrakantisinha@yahoo.com

### Assistant Editors

**Dr. Prateek Tripathi**

Molecular & Computational Biology  
University of Southern California  
TRF 108, 3430 S Vermont Ave  
Los Angeles CA-90089  
Email ; tprateek@dornsife.usc.edu

**Dr. Vishva Deepak Tripathi**

Research Associate  
NRC for Agroforestry,  
Near Pahuj Dam  
Gwalior Road, Jhansi -284 003, U.P.

**Dr. S. B. Yadav**

Department of Agromet  
A A U, Anand, Gujarat,  
e-mail: sbmet@yahoo.in

**Dr. R. P. Kaushal Asstt. Prof.**

Deptt. of Statistics & Social Science  
MSKJUA&T, Banda - 210001 (UP)  
Email: kaushal\_rajendra@india.com

### Executive Editor

**Prof S. K. Tripathi**

Department of Water Resources De-  
velopment and Management  
I.I.T., Roorki-247667,  
e-mail: sankufwt@gmail.com  
sankufwt@iitr.ernet.in

### Coordinating Editor

**Dr. Arvind Kumar Srivastava,**

Dept of Meteorology,  
N D University of Agri. and  
Tech., Faizabad, U P.  
E: arvindmausam@gmail.com

### Technical Editor

**Dr. A. K. Srivastava, Principal**  
Scientist (Agromet)

I I S R Lucknow  
E: arunkumarsrivastava2@gmail.com

### Publication Editor

*Dr. Manisha Verma, B. Sc., B. Tech.*

**Publication Editor (Chief Executive Director)**

**Academic And Research Publications**

H. Office: EC 41, Maya Enclave, New Delhi -110064

Email : manisha\_npp@yahoo.com , [www.manishanpp.com](http://www.manishanpp.com)

© Journal on Agricultural Science and Technology. All rights reserved. No portion of material can be reproduced in part or full without the prior permission of the Editor.

Note : The views expressed herein are the opinions of contributors and do not reflect the stated policies of the Academic And Research Publications. Correspondence: All enquiries, editorial, business and any other, may be addressed to: The Editor-in-chief, International Journal of Agricultural Science and Technology (IJAgST), H. Office: EC 41, Maya Enclave, New Delhi -110064 (INDIA)

Email : manik.sinha@ymail.com; manisha\_npp@yahoo.com, [www.manishanpp.com](http://www.manishanpp.com).

# **International Journal Of Agricultural Science & Technology**

## **Members of Editorial Board**

### **Dr. V. N. Pathak**

Asstt. Prof., Deptt of GPB  
S.M.M. Town College, Ballia, U.P.  
E: vijayanand.pathak76@gmail.com

### **Dr. Alok Kumar Singh**

Deptt of Crop Physiology  
N. D. U A. & T, FAIZABAD-224 229  
.e-mail: aloksingh.agri@gmail.com

### **Dr. B. P. Shahi**

#### **Asstt Prof./SMS (Horticulture)**

Krishi Vigyan Kendra, Meerut.  
e-mail: bhshahi1975@yahoo.com

### **Dr. P. K. Singh**

India Meteorological Department  
Mausam Bhavan, Lodhi Road  
New Delhi-110003, India.  
e-mail: pksingh66@gmail.com

### **Prof J. P. Srivastava**

Head, Deptt of Plant Physiology  
Institute of Agricultural Sciences  
Banaras Hindu University, Varanasi,

### **Dr. Vyas Pandey**

Prof. & Head, Deptt of Agromet  
A A U, Anand, Gujarat,  
e-mail: pandey04@yahoo.com

### **Dr. Akhilesh Srivastava**

Assoc. Prof., Horticulture  
M P U of Agril. & Tech., Udaipur,

### **Dr. Anil Kumar Singh**

Asstt. Prof., Agronomy  
N.D.U.A.& T., Faizabad.  
e-mail: aksmausam@gmail.com

### **Dr. Neeraj Kumar**

Astt Prof, Deptt of Soil Science,  
N. D. U A. & T, FAIZABAD - 224 229  
e-mail: nksoilchem@gmail.com

# International Journal Of Agricultural Science & Technology

Volume No. 3

October 2014

Issue No. 2, 2014

## C o n t e n t s

S. No.	Title	Page No.
1.	Allelopathic effect of <i>Eucalyptus tereticornis</i> clone3 on germination and growth of <i>Vigna mungo</i> L <b>V. D. Tripathi, A. Venkatesh and S. K. Dhyani</b>	1
2.	Use of Medicinal plants for treatment of cancer-A review <b>Malika Pal</b>	11
3.	Effect Of Dairy Effluents On Early Seedling Growth Of <i>Pennisetum Typhoides</i> (L) <b>Goutam Bhakta</b>	17
4.	Study The Effect Of Dairy Effluents On Seed Germination Of High Nutrient Quality Pearl Millet- <i>Pennisetum Typhoides</i> (L.) <b>Goutam Bhakta</b>	22







# ALLELOPATHIC EFFECT OF *Eucalyptus tereticornis* CLONE 3 ON GERMINATION AND GROWTH OF *Vigna mungo* L

V. D. Tripathi, A. Venkatesh and S. K. Dhyani

Central Agroforestry Research Institute, Jhansi, 284 003

Email : [deepakvishva2003@gmail.com](mailto:deepakvishva2003@gmail.com)

(Date of Receipt :24-08-2014 ;

Date of Acceptance for Publication : 28-09-2014)

## Abstract

The study was conducted to determine the allelopathic effect of *Eucalyptus tereticornis* clone 3 on urd (*Vigna mungo* L. Hepper) crop. Leaf extracts of *Eucalyptus tereticornis* clone 3 were used to run the experiment under laboratory condition. The aqueous extracts of leaf showed significant inhibitory effect on germination, root and shoot elongation of urd plants. The inhibitory effect was proportional to the concentrations of the extracts and the higher concentration (15%) had the strongest inhibitory effect. In plant part Leaf extract was more inhibitory effect compared to twig and root extract. From the obtained results, it can be concluded that, eucalyptus seem to be a potential threat to the pulses industry under small-scale farming condition. Therefore, it could be recommend that different remedial practices (like removal of excess leaf litters, planting after the rains) should be done before sowing pulses, in land previously planted with Eucalyptus in order to reduce the potential risks.

**Key words:** Allelopathy, Clone , *Eucalyptus tereticornis*, Germination and *Vigna mungo*.

Pages :10

References: 29

## INTRODUCTION

Agroforestry is a dynamic, ecologically based natural resources management system that, through integration of trees on farms and agricultural landscapes, diversifies and sustains production for increased social, economic and environmental benefits for land users at all levels.

*Eucalyptus* spp. is indigenous to Austral-

ia and has been introduced into many countries, owing to their fast growth and their rising demand for paper and plywood (Cossalter and Pye-Smith 2003, Turnbull 1999). To fill the widening gap between the supply and demand of forest raw materials, many *Eucalyptus* species are even grown in agricultural fields with crops (Lisanetwork and Michelsem 1993, Malik 2004) owing to its fast growth (Cossalter and Pye-Smith 2003) wider adaptability (Gandner 2007,





Johansson and Tuomela 1996) and high productivity (Singh and Toky 1995). The genus *Eucalyptus* predominates tree-planting practices among smallholder farmers in India. The genus is introduced to the country over a century ago, but it is showing an alarming expansion throughout rural India in recent decades. *Eucalyptus* is preferred over other species due to a number of merits that address the need of the farmers. Most farmers describe it as “life saviour”, “safety net” or “tree bank” as it is converted easily and quickly to cash wherever needed. However, some public reactions against planting of *eucalyptus*. The criticisms are based on a range of technical, ecological and socio-economic arguments (Deml Teketay 2000). One of the ecological arguments is that, eucalypts threaten biodiversity and habitat quality by a phenomenon called allelopathy (El Darier 2002). Allelopathy can simply be understood as the ability of plants to inhibit or stimulate growth of other plants in the environment by exuding chemicals.

Many studies have evaluated the allelopathic effects of *Eucalyptus* species (Bajwa and Nazi 2005, Del Moral and Muller 1969, El-Khawas and Shehata 2005). Certain phenolic acids and volatile oils released from the leaves, bark and roots of certain *Eucalyptus* spp. act as allelopathic agents and are harmful to other plant species (Florentine and Fox 2003, Sasikumar *et. al.*, 2002). Chemically allelochemical compounds have open chain molecular structures. These are secondary metabolites that have role in plant-plant, plant-soil, plant-disease, plant-insect and plant predator interactions that may be beneficial or detri-

mental to plant (Tang *et. al.*, 1989). The chemicals have harmful effects on the crop in the eco-system resulting in the reduction and delaying of germination, mortality of seedlings and reduction in growth and yield (Herro and Callaway 2003).

Urd (*Vigna mungo* L. Hepper) is one of the important pulse crops in agroforestry system because of its short duration, protein riched and nitrogen fixing character. The main characteristics of urd i.e reducing fertilizer needs, improving soil structure and providing plant protein are particularly important for developing countries where agricultural production is often limited due to the lack of supply of N fertilizer to the agricultural field.

## MATERIALS AND METHODS

The experiment was conducted in the general laboratory having an average temperature range of 30-32 °C. The CAFRI is located at about 24° 11' N latitude and 78° 17' E longitude and an elevation of 271 meters above sea level (msl). Mean annual rainfall is 960 mm with an average of 52 rainy days per year. Mean maximum temperature ranges from 47.4 °C (June) to 23.5 °C (January) and mean minimum temperature from 27.2 °C (June) to 4.1 °C (December). The main soil types in the region are red (Alfisol) and black (Vertisol).

Seeds of urd (*Vigna radiate* var. PU35, family: leguminaceae) were used. The seeds were thoroughly washed with distilled water and surface sterilized with mucuric chloride for 2 minutes

and rinsed four times with distilled water.

The *Eucalyptus tereticornis* Clone3 water extract were prepared as follows. Fifty, hundred and one hundred fifty grams of air-dried *Eucalyptus tereticornis* clones 3 plant parts viz., leaf, twig and root powder were separately soaked in one litre of distilled water for about 24 hours at room temperature. The mixture were filtered through a double layer muslin cloth to obtain 5%, 10% and 15% concentrations of different plant parts. A complete randomised design was used for this experiment. This experiment was a laboratory bioassay. Four types of extracts from different parts of *E. tereticornis* clones were prepared. Twenty seeds of urd were arranged in 9 cm diameter Petri-dishes on two layers of filter papers under normal laboratory conditions. The experiment was replicated five times with completely

randomised design.

For laboratory bioassay, the germination percentage recorded 7<sup>th</sup> day after sowing and plumule length and radicle length were recorded after 15 day. For measurement of the plumule and radicle length five representative germinated seeds were considered randomly from the twenty seeds in each Petri-dish.

The vigour index was calculated by using following formula given by Abdul Baki and Anderson, (1973).

Vigour index = Germination % X (Root length + Shoot length)

The dickson quality index was calculated by using following formula (Dickson *et. al.*, 1960)

$$\text{Dickson Quality Index (DQI)} = \frac{\text{Total dry weight (gm)}}{\text{Plant height (cm)} + \text{Shoot dry weight (gm)}} + \frac{\text{Collar diameter (mm)}}{\text{Root dry weight (gm)}}$$

Data of germination and growth parameters were subjected to ANOVA using SYSTAT statistical program. Comparison of the mean was performed using Least Significant Difference (LSD) with the control.

## RESULT AND DISCUSSION

### Germination (%)

The result presented in table 1, showed that germination efficiency of urd plants decreased significantly by the 15% leaf extract. Ten percent leaf extract had the next more pronounced negative impact on germination of urd, which is significantly higher as compared to 5% leaf extract.

### Vigour index:

The vigour index was presented in table no. 2. The result revealed that maximum inhibition was recorded by the 15% leaf

extract, which is significantly lower at 10%, and 5% leaf extract. Root extract showed less inhibition at 5% concentration. Control (0%) had higher vigour index as compared with plant parts and concentrations.

### Shoot length (cm)

The shoot length of urd was highly suppressed by the 15% leaf extract. The maximum shoot length was recorded by 5% root extract. Control (0%) showed maximum shoot length (Table 3).

### Root length (cm)

The root length of urd was significantly inhibited by 15% leaf extract. However, maximum root length was found in 5% twig extract. In plant parts, maximum inhibition was observed by leaf followed by root and twig (Table 4).

### Shoot and Root Dry Weight (gm/5 seedlings)

Shoot dry weight of urd was significantly superior in 0% (control) (95.9 gm/5 seedlings). The 15% concentration (42.97 gm/5 seedlings) of leaf extract was recorded maximum inhibition effect followed by 10% (56.1 gm/5 seedlings) and 5% (64.0 gm/5 seedlings). The maximum root dry weight was observed in control (60.7 gm/5 seedlings). The significantly inhibition on root dry weight was observed by 15% concentration (18.2 gm/5 seedlings) followed by 10% (22.8 gm/5 seedlings) and 5% (27.3 gm/5 seedlings) (Fig.1). In





plant parts of *E. tereticornis* clones3, maximum suppression in shoot dry weight was found by leaf followed by twig and root. While, root dry weight was inhibited by leaf followed by root and twig (Fig. 2).

### Dickson quality index (DQI)

In Dickson quality index maximum suppressing was recorded by leaf extract of 15% concentration followed by root extract and twig extract at 15% concentration. Significantly, higher DQI was observed in control (0%). It was observed that 10% of twig extract take maximum DQI followed by leaf and root extract (Fig. 3).

*Eucalyptus* monoculture plantations are reported to support either very little or almost negligible under-storey vegetation (Del Moral and Muller 1969, Singh *et. al.*, 1993). The species diversity index is also highly reduced under eucalypt monoculture plantations when compared with the other native plantations. Allelopathy has often been considered as a possible reason for the species depletion (Suresh and Rai 1987, Kohli *et. al.*, 1992). Venkatesh and Tripathi (2010) have reported significant reduction in the density, root and shoot length, biomass, and economic yield of forage crops under *E. tereticornis* clones. *Eucalyptus* are reported to release a number of volatile and non-volatile allelochemicals that affect growth of the associated vegetation (Kohli 1990). Various volatile terpenes like limonene, cineole,

citronellal, citronellol,  $\alpha$ -pinene, and grandinol, etc. identified from the crude oil are highly toxic and affect the germination and growth of native vegetation (Baker 1966, del Moral and Muller 1970, Kohli *et. al.*, 1992). A number of phenolic compounds viz., coffecic, coumaric, ferric, gallic, gen-tisic, hydroxybenzoic, syringic and vanillic acids and catechol in bark, fresh leaves, root and seed leachates of *Eucalyptus* spp. were identified in the soil and leaves of *Eucalyptus* (Kohli and Singh 1991). Similar inhibitory mechanism of *Eucalyptus* spp. on test crops were already reported by Suresh and Rai (1987) and Sasikumar *et. al.*, (2001). In addition, the leachates and extracts from the eucalypt leaves, litter, bark, flowers, and leaf mulch have been reported to reduce the germination and initial growth of a number of plant species (Singh and Bawa 1982, Sidhu and Hans 1988 and Kohli 1990).

### CONCLUSION

This study was carried out to investigate the Phyto-toxicity of *Eucalyptus tereticornis* clones3 on germination and growth of urd (*Vigna mungo* L. Hepper). Germination and all the growth parameters of urd was reduced significantly under 15% leaf aqueous extract of clone3 of *Eucalyptus tereticornis*. Among the plant parts, the maximum toxicity was exhibited by leaf. Toxicity of the extract was in the order of leaves > twigs > roots and the concentration was in the order of 15% > 10% > 5% > 0%.

Table 1. Germination percentage in urd plants after 7 days of sowing affected by *E. tereticornis* clone3 extract.

Extract Source (P)	Extract Concentration (C)				
	0%	5%	10%	15%	Mean
Leaf	85	38	24	9	39
Twig	85	56	50	41	58
root	85	61	55	47	62
Mean	85	51.67	43.00	32.33	
	P	C	PxC		
LSD <sub>0.05</sub>	1.77	2.05	3.55		

Table 2. Vigour Index of urd plants affected by *E. tereticornis* clone3 extract.

Extract Source (P)	Extract Concentration (C)				
	0%	5%	10%	15%	Mean
Leaf	2783	293	98	18	798
Twig	2783	1223	881	415	1325
root	2783	1406	1137	664	1498
Mean	2783	974	705.33	365.67	
	P	C	PxC		
LSD <sub>0.05</sub>	51.87	59.90	103.74		

Table 3. Shoot length (cm) of urd plants affected by *E. tereticornis* clone3 extract.

Extract Source (P)	Extract Concentration (C)				
	0%	5%	10%	15%	Mean
Leaf	17.1	5.6	3.4	1.4	6.9
Twig	17.1	10.3	5.6	5.7	10.1
root	17.1	11.7	11.0	7.3	9.6
Mean	17.1	9.20	6.67	4.8	
	P	C	PxC		
LSD <sub>0.05</sub>	0.39	0.45	0.78		

Table 4. Root length (cm) of urd plants affected by *E. tereticornis* clone3 extract.

Extract Source (P)	Extract Concentration (C)				
	0%	5%	10%	15%	Mean
Leaf	11.1	2.0	2.1	0.9	4.0
Twig	11.1	7.6	7.3	4.1	7.5
root	11.1	6.7	5.9	5.1	6.2
Mean	11.1	5.43	5.1	3.37	
	P	C	PxC		
LSD <sub>0.05</sub>	0.20	0.24	0.41		




Fig. 1. Shoot dry weight and root dry weight of urd plants affecting by *E. tereticornis* clone3 different extract concentrations.




Fig. 2. Shoot dry weight and root dry weight of urd plants affecting by *E. tereticornis* clone3 plant parts extract.




Fig. 3. Dickson Quality Index of urd plants affecting by *E. tereticornis* clone3 extract.





## REFERENCES

1. Abdul Baki, A.A. and Anderson, J.D. **1973**. Vigour determination in soybean seed by multiple criteria. *Crop Sci.* 13: 360-363.
2. Bajwa, R. and Nazi, I. **2005**. Allelopathic effects of *Eucalyptus citriodora* on growth, nodulation and AM colonization of *Vigna radiata* (L) Wilczek. *Allelopathy Journal* 15: 237-246.
3. Baker, H.G. **1966**. Volatile growth inhibitors from *Eucalyptus globulus*. *Madrono*. 18: 207-210.
4. Cossalter, C. and Pye Smith C. **2003**. Fast-wood Forestry: Myths and Realities. Center for International Forestry Research, Indonesia. pp 9.
5. Del Moral, R. and Muller, C.H. **1970**. The allelopathic effects of *Eucalyptus camaldulensis*. *Amer. Midl. Nat.* 83: 254-282.
6. Del Moral, R. and Muller, C.H. **1969**. Fog drip: a mechanism of toxin transport from *Eucalyptus globulus*. *Bull. Torrey Bot. Club* 96: 467-475.
7. Deml, Teketay. **2000**. The ecological effects of eucalyptus: Guide for making wise an informed decision. In: Kinf Abebe (eds.), *The Eucalyptus Dilemma: The Socio-economic aspects of eucalyptus*, Addis Ababa. pp. 1-45.
8. Dickson, A. Leaf, A.L. and Hosner, J.F. **1960**. Quality appraisal of white spruce and white pine seedling stock in nurseries. *For. Chron.* 36: 10-13.
9. El Darier, S.M. **2002**. Allelopathic effect of *Eucalyptus rostrata* on growth, nutrient uptake and metabolite accumulation of *Vicia faba* L. and *Zea mays* L. *Pakistan Journal of Biological Sciences* 5(1): 6-11.
10. El Khawas, S.A. and Shehata, M.M. **2005**. The Allelopathic potentialities of *Acacia nilotica* and *Eucalyptus rostrata* on monocot (*Zea mays* L.) and dicot (*Phaseolus vulgaris* L.) plants. *Biotechnology* 4: 23-34.
11. Florentine, S.K. and Fox, J.E.D. **2003**. Allelopathic effects of *Eucalyptus victrix* L. on *Eucalyptus* species and grasses. *Allelopathy Journal* 11: 77-83.
12. Gardner, R.A.W. **2007**. Investigating the environmental adaptability of promising subtropical and cold tolerant *eucalyptus* species in warm temperate climate zone of KwaZulu-Natal, South Africa. *Southern Hemisphere Forestry Journal* 69:27-38.
13. Herro, J.L. and Callaway, R.M. **2003**. Allelopathy and exotic plant invasion. *Plant and Soil* 256: 29-39.
14. Johansson, S. and Tuomela, K. **1996**. Growth of 16 provenances



- of *Eucalyptus microtheca* in a regularly irrigated plantation in eastern Kenya. *Forest Ecology and Management* 82: 11-18.
15. Kohli, R.K. and Singh, D. **1991**. Allelopathic impact of volatile components from *Eucalyptus*. on crop plants. *Biol. Plant.* 33: 475–483.
  16. Kohli, R.K., Singh, D. and Rani, D. **1992**. Allelopathic potential of volatile terpenes of *Eucalyptus*. In: Environmental Concerns in Rights-of-Way Management. pp. 102–106.
  17. Kohli, R.K. **1990**. Allelopathic Potential of *Eucalyptus*. Project Report MAB-DOEn. Project, India.
  18. Lisanework, N. and Michelsen, A. **1993**. Allelopathy in agroforestry systems - the effects of leaf extracts of *Cupressus lusitanica* and three *Eucalyptus*. spp. on four Ethiopian crops. *Agroforestry System* 21: 63- 74.
  19. Malik, M.S. **2004**. Effects of aqueous leaf extracts of *Eucalyptus globulus* on germination and seedling growth of potato, maize and bean. *Allelopathy Journal* 14: 213-219.
  20. Sasikumar, K., Vijayalakshmi, C. and Parthiban, K.T. **2001**. Allelopathic effects of four Eucalyptus species on redgram (*Cajanus cajan, L.*). *Journal of Tropical Agriculture* 39(2): 134-138.
  21. Sasikumar, K., Vijayalakshmi, C. and Parthiban, K.T. **2002**. Allelopathic effects of Eucalyptus on blackgram (*Phaseolus mungo L.*). *Allelopathy Journal* 9: 205-214.
  22. Sidhu, D.S. and Hans, A.S. **1988**. Preliminary studies on the effect of *Eucalyptus* leaf-litter on accumulation of biomass in wheat. *Journal of Tropical For.* 4: 328–333.
  23. Singh, D. Kohli, R.K. and Jerath, N. **1993**. Impact of *Eucalyptus* and other plantations on phytodiversity in India. In: Proc. Int. Conf. Forest Vegetation Management-Ecology, Practice and Policy. Gjerstad D.H., Ed., USA. pp. 152–159.
  24. Singh, R. and Bawa, R. **1982**. Effects of leaf leachates from *Eucalyptus globulus* Labill. and *Aesculus indica* Colebr. on germination of *Glaucium flavum* Crantz. *Ind. J. Ecol.* 9: 21–28.
  25. Singh, V. and Toky, O.P. **1995**. Biomass and net primary productivity in *Leucaena*, *Acacia* and *Eucalyptus*, short-rotation, high-density (Energy) plantations in arid India. *Journal of Arid Environments* 31: 301-309.
  26. Suresh, K.K. and Vinaya, Rai R.S. **1987**. Studies on the allelopathic effects of some agroforestry tree crops. *International Tree Crops Journal.* 4(2-3): 109-115.
  27. Tang, C.S., Komai, K. and



- Haung, R.S. **1989**. In Phytochemical Ecology: allelochemicals, mycotoxins and insect pheromones and allomones. CH Chou and GR Waller (Eds.). *Institute of Botany Academia Sinica Monograph series 9, Taipei, Roc*, pp. 217-223.
28. Turnbull, J.W. **1999**. *Eucalyptus*-plantations. *New Forest* 17:37-52.
29. Venkatesh, A. and Tripathi, V.D. **2010**. Phyto-toxicity of *Eucalyptus tereticornis* clones on forage crops. *Range Management and Agroforestry. Symposium Issue (B)*: 132-133.







## USE OF MEDICINAL PLANTS FOR TREATMENT OF CANCER-A Review

**Malika Pal**

Principal  
Extol Institute of management  
Bhopal

Email : [drmalikapal@gmail.com](mailto:drmalikapal@gmail.com)

(Date of Receipt :27-08-2014 ;

Date of Acceptance for Publication : 06-09-2014)

### Abstract

Cancer is a major public health problem in both developed and developing countries. It was estimated that there were 10.9 million new cases, 6.7 million deaths, and 24.6 million persons living with cancer around the world in 2002. <sup>(15,16)</sup> And the number of cancer deaths are increasing, because most of the times the intervention is too late because patients put up with the symptoms instead of seeking treatment. <sup>(4)</sup>

**Key Words:** Cancer, Symptoms, Treatment, Cause of Cancer or The Molecular Basis of Cancer, Medicinal Plants And Traditional Medicine.

Pages: 6

References: 25

### INTRODUCTION

Cancer is a fascinating biological problem and can be defined as a disease involving heritable defects in cellular control mechanism resulting in the formation of malignant and usually invasive tumors. A cancer arises from a single cell that undergoes a permanent change such that it goes on multiplying defying the architectural requirements of the tissue, the organ and the organism. The cancer cells

morphologically and metabolically differs from normal cells of the body. Their source of energy is predominantly anaerobic. The tumor cells are primarily concerned with growth, not function. It acquires nitrogenous building blocks from the body stores to satisfy the continual demand for protein synthesis.

The cause of cancer or the molecular basis of cancer can be traced to, a non le-

thal damage to the chromosomes. Such genetic damage may be acquired by the action of environmental agents such as chemicals, radiations, viruses or it may be inherited in germ line<sup>(1)</sup>. Free radicals are molecules produced when the body breaks down food or by environmental exposures like smoke, radiation and tobacco. These free radicals can damage cells and play an important role in leading to heart disease, and cancer<sup>(2)</sup>.

Antioxidants are the substance that protects the cells against the harmful effects of free radicals. They are present in foods and prevent the oxidation, a process which causes damage to the body. When the body uses oxygen, the by products which are produced are called free radicals which leads to cell damage. Antioxidants act as free radical scavengers and they prevent or repair the damage caused by the free radicals. Antioxidants are not produced by the body; they have to be taken through food. Plants are good source of antioxidants<sup>(3,5)</sup>.

Hippocrates, father of modern medicine stated "Your food shall be your medicine and your medicine shall be your food." Concurrent nutritional deficiency plays a fundamental role in genesis of tumor and suggests that the disease process could be reversed by re-supplying the required nutrients in adequate amount.<sup>(6)</sup> Their holistic approach puts in place a balanced whole food diet, properly assessed mineral and vitamin supplementation and the usual herbal support for the organs of elimination, especially the liver, using plants such as dandelion root (*Taraxacum officinale* radix), milk thistle (*Carduus marianus*) and artichoke juice (*Cynara scolymus*).<sup>(7)</sup> Traditionally many plants and their

parts have been used for the treatment of cancer or the symptoms arising due to it. Plants like - *Vinca rosea*, *Allium sativum* were frequently used as anti-inflammatory and anti tumor agents. Modern natural holistic approach to treat cancer has also accepted the importance of traditional knowledge. In Philippines "banaba" (*Lagerstroemia speciosa*) and "tsaang gubat" (*Carmona retusa*) has tested positive as potent anti cancer drugs<sup>(4)</sup>. In India 45000 species of plants are known, out of which 7000-7500 plants have been listed as plants of medicinal value. The ayurvedic system uses 700 plants, siddha system uses 600 and unani system uses 700 plants approximately, for medicinal purposes. Plants like *Abrus precatorius* (Ghungchi), *Acorus calamus*, *Ageratum conyzoides* L. (Asteraceae), *Aglaia roxburgiana* (Priyangu), *Alnus japonica*, *Bauhinia variegata*, *Cassia fistula*, *Catharanthus roseus* (*Vinca rosea*, Sadabahar), *Carduus marianus*, *Crocus sativus* (Saffron), *Curcuma longa* L., *Cynara scolymus*, *Ervatania heyneana*, *Hygrophila spinosa*, *Hippocratea murcantha*, *Indigofera mysorensis*, *Ocimum sanctum* (Tulsi), *Olea polygama*, *Phyllanthus amarus*, *Plumbago rosea*, *Podophyllum hexandrum*, *Semecarpus anacardium*, *Solenum dulcamara*, *Solenum indicum*, *Solenum khasianum*, *Solenum suratanse*, *Taraxacum officinale* radix, *Terminalia arjuna*, *Terminalia chebula*, *Trigonella foenumgraecum*, *Venda parviflora*, *Wedelia calendulacea*, *Withania somnifera* & *Zingiber capitatum* are still used traditionally as herbal drugs against various types of tumors such as sarcoma, lymphoma, carcinoma and leukemia.<sup>(7,8,9,17,19,20)</sup> Medicinal plants contains anti-oxidants eg:- vitamins (A, C, E, K), carotenoids, flavinoids (flavones, isoflavons, flavonones, anthocyanins, catechins, isocatechins), polyphenoles (ellagic acid, gallic acid,



tannins), saponins, enzymes and minerals (selenium, copper, manganese, zinc, chromium, iodine).<sup>(10)</sup> Several mechanisms have been proposed to explain the cancer-preventive effects of plants. These include inhibition of mutagenesis by inhibiting the metabolism, inhibition of DNA adduct formation, free-radical scavenging, and effects on cell proliferation and tumor growth.<sup>(24)</sup>

The interest in traditional medicine is not new, in 1925, Chatterji successfully used a copper margosate-ester in patients with head and neck cancer. Head and neck cancer constitutes a major problem in India. Hence, any potentially complementary therapy is worth exploring in these patients, where it could enhance the quality of their lives.<sup>(11)</sup> The wound-healing, anti-inflammatory and antimutagenic activities of turmeric have been demonstrated convincingly.<sup>(12)</sup> Hastak *et. al.*, have shown its beneficial effect in oral submucous fibrosis - a precancerous condition<sup>(13)</sup>. In dimethyl benzantracene (DMBA)-induced experimental breast cancer, curcumin has shown a significant reduction in carcinogenesis.<sup>(14)</sup> *Terminalia chebula* was active against leukemia cell lines. *Terminalia chebula*, *Acorus calamus*, *Bauhinia variegeta* and *Phyllanthus amarus* were found to be effective against prostrate D U 145 cell lines<sup>(9)</sup> *Curcuma longa* L.(Zingiberaceae), which exhibit anti-inflammatory, anti-human immunodeficiency virus, anti-bacteria, antioxidant effects and nematocidal activities. Curcumin is a major component in *Curcuma longa* L., being responsible for its biological actions. In vitro, curcumin exhibits anti-parasitic, antispasmodic, anti-inflammatory and gastrointestinal effects; and also inhibits carcinogenesis and cancer growth.<sup>(19)</sup> The stem bark of the

Betulaceae plant *Alnus japonica*, which is indigenous to Korea, has been used as a popular folk medicine for hepatitis and cancer.<sup>(20)</sup>

With nature as a primary source and inspiration three basic strategies, are being followed for treatment of cancer. The first strategy in the traditional approach i.e. use of traditional system of treatment in which a holistic approach is followed. Natural cancer therapy focuses on eliminating the cancer cells but also focuses on stopping the process that initiated the cancer growth. Plants cleanse and revitalize our bodies from the inside. Use of traditional low fat, high fiber diet, rose oil, blackstrap molasses etc, is suggested with greater stress on fresh fruits vegetables and non refined products.<sup>(6)</sup>

The second strategy focuses on blending of the benefits of both the systems. In this line of treatment the main treatment is done using modern surgical processes along the use of chemotherapy etc but botanicals are given to counteract the harmful side effects of these powerful drugs. For example plant *Tinospora cordifolia* Hook ( Guduchi) was used as a standardized formulation before and after cancer chemotherapy to reduce the incidence of nausea, vomiting and granulocytopenia.<sup>(18)</sup> The therapy commonly called as phytotherapy uses simple cost-effective measures that can provide greater patient comfort, more effective treatment of the patient within the hospital and, it is hoped, a longer, more fulfilling survival period.<sup>(7)</sup> But the third strategy which involves identification, isolation and characterization of various bioactive compounds present in medicinal plants.<sup>(2, 19, 20)</sup> is the most favoured one. The stress is on clinically us-





ing secondary metabolites and their derivatives to combat cancer. <sup>(21)</sup>

Currently, over 50% of drugs used in clinical trials for anticancer activity were isolated from natural sources or are related to them. <sup>(22)</sup> A number of active compounds have been shown to possess anticancer activity; these include flavonoids, diterpenoids, triterpenoids, and alkaloids. <sup>(23)</sup> The bioactive compound responsible for antiradical activity of *A. conyzoides* is <sup>3</sup>, <sup>5</sup>, <sup>7</sup>, <sup>4</sup>-tetrahydroxyflavone (kaempferol)., <sup>(25)</sup> Curcumin in *Curcuma longa* L.(Zingiberaceae)<sup>(19)</sup>.

Now it is clear that the diseases, caused by free radicals, could be managed by enhancing anti-oxidant defense systems. The discovery of new enzymes inhibitors and antioxidant compounds from medicinal plants would surely help overcome the cancer agents. As preparation of standardize dose & dosage regime which are very crucial for the treatment will become easier after isolation of the compound. This will enable us to produce the bioactive compound or its chemical derivatives en masse. So the future of cancer treatment lies in the hands of reverse pharmaceuticals that identify the natural compounds in plants and synthesize their chemical derivatives.

## ACKNOWLEDGEMENT

The authors are grateful to the authors / editors of all those articles, journals and books from where the matter for this has been reviewed and discussed.

## REFERENCES

1. Neoplasia-Cell Biology Fundamentals and applications, M.L. Gupta and M.L. Jangir., *Agrobios* **2001**.634-652.
2. Medicinal plants may help treat cancer : Experts say – 13 Jan **2004**., Dawn the INTERNET Edition , [http:// DAWN.com](http://DAWN.com) staff reporter.
3. Antioxidants- <http://www.health-caremagic.com/health> Resource.
4. Medicinal plants vs cancer. Henrylito D. Tacio. Sept 19-**2005**. <http://sunstar.com.ph/e-magazine/index.htm>.
5. Understanding free radicals and antioxidants. <http://www.healthcheck-system.com/info.htm>.
6. Cancer: a single disease with multitude manifestations? Peter Grandics. *J. of Carcinogenesis*. **2003**,2-9.
7. Can Medicinal Plants Beat Cancer? Keith Stelling, MNIMH, Dip-Phyt alive #222, April **2001**.
8. Ethano medicinal plants for prevention and treatment of tumors. Sharma and Pandey. *International journal of green pharmacy* march **2009**. 2-5.
9. Gaidhani SN, Lavekar GS, Juvekar AS, Sen S, Singh A, Kumari S. In-vitro anticancer activity of standard extracts used in ayurveda. *Phcog Mag* **2009**;5:425-9.



10. Gupta V K, Sharma S K. plants as naturals anti- oxidants. Nat. Pro. Rad. 2006, 326-34. Kaur C, Kapoor H C., Anti- oxidant activity and total phenolic content of some Asian vegetables. *Int. J. Food Sci. Tech.* **2002**; 37;153-61)
11. Chatterji KK. A note on margo-sa oil in the treatment of cancer. *Lancet* **1925**;2:1063-4.
12. Chandra D, Gupta SS. Antiin-flammatory and antiarthritic ac-tivity of volatile oil of *Curcuma longa* (haldi). *Indian J Med Res* **1972**;60:138-42.
13. Hastak KA, Lubri N, Jakhi SD, More C, John A, Ghaisas SD, *et. al.*, Effect of turmeric oil and turmeric oleoresin on cytogenetic damage in patients suffering from oral sub-mucous fibrosis. *Cancer Lett* **1997**; 116:265-69. *urcuma longa* (haldi). *Indian J Med Res* 1972;60: 138-42.
14. Bhide SV, Azuine MA, Lahiri M, Telang NT. Chemoprevention of mammary tumor virus-induced and chemical carcinogen-induced rodent mammary tumors by natu-ral plant products. *Breast Cancer Res Treat* **1994**; 30:233-42.
15. Parkin DM, Bray F, Ferlay J, Pisani P. Global cancer statistics, **2002**. CA: *Cancer J Clin* 2005;55:74-108.
16. Shoeb M. Anticancer agents from medicinal plants. *Bangl J Pharmacol* **2006**;1:35-41.
17. Adebayo AH, Tan NH, Akinda-hunsi AA, Zeng GZ, Zhang YM. Anticancer and antiradical scav-enging activity of *Ageratum cony-zoides* L. (Asteraceae). *Phcog Mag* **2010**;6:62-6
18. Late Shringi MS. Personal com-munication **2002**.
19. Araujo,CC, Leon,LL.Biological activities of *Curcuma longa* L. *Phyto-ther Res.* **2001** Jul; 96 (5):7234-8.
20. Sang Tae Km., Jung Do Kim, Seung Hee Ahn, Young Ik Lee, Young Seok Jeog **2004**. Hepato protective & anti oxidant effects of *Alnus japonica* extracts on aceta-minophen induced hepato toxicity in rats.*Phytother Res.* **2004**.Dec 18 (12):971-5.
21. Adebayo AH, Tan NH, Akinda-hunsi AA, Zeng GZ, Zhang YM. An-ticancer and antiradical scaveng-ing activity of *Ageratum conyzoides* L. (Asteraceae). *Phcog Mag* **2010**;6:62-6
22. Newman DJ, Cragg GM. Natu-ral products as sources of new drugs over the last 25years. *J Nat Prod* **2007**;70:461-77.
23. Han HJ, Tan NH, Zeng GZ, Fan JT, Huang HQ, Ji CJ, *et. al.*, Natural



inhibitors of DNA topoisomerase I with cytotoxicities. *Chem Biodivers* **2008**;5:1364-8.

24. Shukla Y, Kalra N. Cancer chemoprevention with garlic and its constituents. *Cancer Lett*

**2007**;247:167-81.

25. Adebayo AH, Tan NH, Akindahunsi AA, Zeng GZ, Zhang YM. Anticancer and antiradical scavenging activity of *Ageratum conyzoides* L. (Asteraceae). *Phcog Mag* **2010**;6:62-6.





## EFFECT OF DAIRY EFFLUENTS ON EARLY SEEDLING GROWTH OF *Pennisetum typhoides* (L)

**Goutam Bhakta**

Rukminipur K.L Vidyapith (High),  
Egra, Purba Medinipur, West Bengal

Email : [ani.ezee@gmail.com](mailto:ani.ezee@gmail.com)

(Date of Receipt :14-09-2014 ;

Date of Acceptance for Publication : 02-10-2014)

### Abstract

With the growing industrialization and urbanization environment degradation has now become challenging global problem. Among the industries food processing industrial effluents released from the dairy industries are rich in various kinds of nutrients like phosphate, Calcium, Nitrogen, Magnesium etc. and has good potential in utilization of released effluents as source of nutrients for the crop plants. To study the effect of dairy effluents on early seedling growth of *Pennisetum typhoides* (L). different parameters like i) Plumule and radicle ratio, ii) Fresh wt. and dry wt of germinated seeds, iii) Height of the stem, iv) Fresh wt. and dry wt. of stem, v) Root length, vi) no of leaves, vii) internodal length etc, of plants under various concentrations of milk plant effluent in compost rich and compost free soil are taken. The vegetative growth of pearl millet increases with the increase in concentration of effluent upto 80 percent decreases when irrigated with 100. conc. of effluent.

**Key Words:** Pearl Millet, Seedling Vegetative Growth, Dairy Effluent.

Pages: 05

References: 11

### INTRODUCTION

Pearl millet, *Pennisetum typhoides* (L) is the basic staple for households in the poorest countries and among poorest people because of having high protein and high fat content Chemical fertilizer and industrialisation cause great hazards to the crop field, but dairy effluents released from milk

plant due to presence of varied groups of chemical compounds including nutrients like phosphate, magnesium, calcium etc. help in soil fertility and would also increase productivity of the land. The study related to effect of dairy effluents on early seedling growth of *Pennisetum typhoides*(L) is of great emphasis.



## MATERIAL AND METHODS

For studying physico-chemical analysis of soil collected from Balsamand area, Hisar air dried soil samples passing through 2.0 mm sieve were used, ph, electrical conductivity, calcium carbonate, organic carbon, nitrogen, phosphorous, potassium estimation are done and for physico-chemical analysis of dairy effluents released from milk plant, sirsa, Haryana, Ph, electrical conductivity, total dissolved solids, calcium, magnesium, sulphate, phosphate, chloride, BOD, COD are done. To carry out the effect dairy effluents on early seedling growth first twenty seeds sterilized in  $HgCl_2$  for 2 minutes were soaked in water in a beaker for 8 hours. Total thirty six earthen pots (18 for compost soil and 18 for without compost each filled with 2.0 kg soil numbering with 1, 2, 3 .....35,36 in each pot numbered 1-15, 100 gms vermicompost was also mixed thoroughly, then these pots were irrigated with different doses of effluent i.e control 20%, 40%, 60%, 80%, and 100% regularly up to 60 days. then different parameters of early seedling growth were observed and recorded minutely.

## RESULTS AND DISCUSSION

**Table 1 :**

The height of stem (cm), under compost and without compost, of pearl millet as influenced by different concentrations of effluent.

Concentrations	Height of Stem (cm)							
	Compost				Without Compost			
	15 <sup>th</sup>	30 <sup>th</sup>	45 <sup>th</sup>	60 <sup>th</sup>	15 <sup>th</sup>	30 <sup>th</sup>	45 <sup>th</sup>	60 <sup>th</sup>
	Day				Day			
Control	4.3	6.2	11.3	40.4	3.4	5.7	9.8	28.5
20%	4.3	5.8	11.8	42.2	3.1	6	10.6	39
40%	4.5	6.3	12.5	45.4	3.5	6.6	11.4	43.8
60%	4.6	6.3	13.1	47.4	4.0	7.0	12.2	47.2
80%	5.2	6.5	13.5	59	4.3	8.8	12.5	52
100%	3.1	4.9	10.7	38.5	3	5.5	8.8	26.7

The stem height of pearl millet was observed to be maximum at 80 % concentration of effluent under both compost (59cm) and without compost (52cm) after 60 days of sowing.

**Table 2**

The fresh weight and dry weight of stem (gm), under compost and without compost, of pearl millet as influenced by different concentrations of effluent.

Concentrations	Stem			
	With Compost		Without Compost	
	Fresh Wt. (gm)	Dry Wt. (gm)	Fresh Wt. (gm)	Dry Wt. (gm)
Control	31	5	12	2.2
20%	36.2	4.2	18.6	2.7
40%	41.6	5	19	2.7
60%	51.6	5.3	21.1	2.9
80%	54	6.8	22.5	3.6
100%	30	4	11.5	2.0

The fresh wt. and dry wt. stem of pearl millet were observed to be maximum at 80% concentration of effluent under both compost (fresh wt =54 gm, dry wt. = 6.8 gm) and without compost (fresh wt = 22.5 gm, dry wt. = 3.6 gm) and were observed to be minimum at 100% conc. of effluent under both compost (fresh wt = 3. gm, dry wt. 4gm) and without compost (fresh wt 11.5 gm, dry wt. = 2.0 gm)

**Table 3 :**

The root length (cm), under compost and without compost, of pearl millet as influenced by different concentrations of effluent, after 60 days of sowing.

The root length was observed to be maximum at 80% concentration of effluent under both with compost (root length 42.4

Concentrations	Root length (cm)	
	Compost	Without Compost
Control	36.8	37.8
20%	38.2	38
40%	40	38.5
60%	41.2	39.2
80%	42.4	40.1
100%	39.6	36.9

cm) and without compost (root length =40.1 cm) and, minimum at control (36.8 cm) under with compost where as in case of without compost the minimum root length (36.9 am) was observed at 100% con of effluent.

**Table 4:**

The fresh weight and dry weight of root (gm), under compost and without compost, of pearl millet as influenced by different concentrations of effluent, after 60 days of sowing.

Concentrations	Root			
	With Compost		Without Compost	
	Fresh Wt. (gm)	Dry Wt. (gm)	Fresh Wt. (gm)	Dry Wt. (gm)
Control	2.2	1.09	1.06	0.57
20%	2.74	1.15	1.11	0.59
40%	3.21	1.17	1.56	0.64
60%	3.7	1.23	1.72	0.8
80%	5.80	1.66	1.80	0.88
100%	2.20	1.03	0.95	0.55

Fresh wt. and dry wt. root were observed to be maximum at 80 percent con. of effluent both compost (fresh = 5.80gm, dry wt. = 1.66 gm) and without compost ( fresh wt. = 1.80 gm. dry wt. =.88), minimum at 100 percent conc. of effluent under both with compost (fresh wt = 2.20 gm. dry wt. = 1.03 gm), without compost ( fresh wt

= .95 gm, dry wt. = .55 gm).

**Table 5:**

The number of leaves, under compost and without compost, of pearl millet as influenced by different concentrations of effluent.

Concentrations	No. of leaves							
	Compost				Without Compost			
	15 <sup>th</sup>	30 <sup>th</sup>	45 <sup>th</sup>	60 <sup>th</sup>	15 <sup>th</sup>	30 <sup>th</sup>	45 <sup>th</sup>	60 <sup>th</sup>
	Day				Day			
Control	4	4	6	8	3	4	5	7
20%	3	4	6	9	3	4	6	8
40%	4	4	7	9	3	4	6	8
60%	4	4	7	10	3	3	6	8
80%	3	4	7	11	3	4	7	9
100%	3	4	6	7	3	4	5	6

Number of leaves was observed to be maximum at 80 percent conc. of effluent under both with compost (11) and without compost (9), minimum at 100 percent conc. of effluent under both with compost (7), without compost (6).

**Table 6 :**

The collar diameter of stem (cm), under compost and without compost, of pearl millet as influenced by different concentrations of effluent.

Concentrations	Collar Diameter (cm)							
	Compost				Without Compost			
	15 <sup>th</sup>	30 <sup>th</sup>	45 <sup>th</sup>	60 <sup>th</sup>	15 <sup>th</sup>	30 <sup>th</sup>	45 <sup>th</sup>	60 <sup>th</sup>
	Day				Day			
Control	1.06	1.60	2.3	4	1.02	1.36	1.77	2.84
20%	1.10	1.61	2.76	3.92	0.84	1.34	1.9	2.6
40%	1.13	1.63	2.75	3.84	0.86	1.38	1.96	2.70
60%	1.2	1.60	2.74	3.77	0.8	1.38	1.95	2.58
80%	0.93	1.34	2.70	3.70	0.88	1.4	2	2.51
100%	0.95	1.33	2.65	3	0.97	1.43	1.6	2.36

Collar diameter of stem observed to be maximum at control treatment under both with compost (4cm) and without compost (2.84 cm) after 60days of sow-

ing, minimum at 100% conc. of effluent both with compost (3.0cm) without compost (2.36cm) after 60 days of sowing.

### 7.(a)

Internodal lengths (cm) of pearl millet as influenced by different concentrations of effluent, after 60 days of sowing (**compost**).

Concentration	Internodal length (cm) (Compost)									
	1 <sup>st</sup>	2 <sup>nd</sup>	3 <sup>rd</sup>	4 <sup>th</sup>	5 <sup>th</sup>	6 <sup>th</sup>	7 <sup>th</sup>	8 <sup>th</sup>	9 <sup>th</sup>	10 <sup>th</sup>
Control	4.34	6.5	9.34	8.67	8.2	6.9	6.2	4.2		
20%	1.93	4.3	5.40	6.65	5.86	4.58	2.53	0.8	2.2	
40%	3.63	8.53	9.6	9.25	8.84	7.92	6.06	2.32	1.8	
60%	1.97	6.46	7.70	7.65	7.59	6.44	5.04	4.2	0.5	
80%	4.07	8.66	10.28	10.45	9.10	8.16	7.38	4.06	2.4	1.8
100%	5.20	8.24	8.16	8.04	8.17	6.70	5.2	4.2		

### 7.(b)

Internodal lengths (cm) of pearl millet as influenced by different concentrations of effluent, after 60 days of sowing (**without compost**).

Concentration	Internodal length (cm) (Without Compost)									
	1 <sup>st</sup>	2 <sup>nd</sup>	3 <sup>rd</sup>	4 <sup>th</sup>	5 <sup>th</sup>	6 <sup>th</sup>	7 <sup>th</sup>	8 <sup>th</sup>	9 <sup>th</sup>	10 <sup>th</sup>
Control	2.36	3.28	4.06	4.2	4.54	3.2	5.94			
20%	1.5	6.12	7.04	7.18	7.39	6.4	4.09	5.34	2.6	
40%	3.82	6.14	8.3	8.16	5.05	4.2	5.72	3.6	2.3	
60%	5.44	6.86	7.16	8.14	9.35	8.36	8.16	7.1		
80%	3.96	7.95	8.72	10.44	9.59	6.8	6.34	2.2	8.2	
100%	1.66	3.90	4.14	6.69	4.99	2.75	2.2			

In case of compost, the minimum internodal length was recorded 20 percent conc. of effluent and in case of without compost minimum internodal length was recorded with control treatment where as maximum at 80% conc. of effluent under both compost and without compost.

## CONCLUSION

Pearl millet crop irrigated with 100 percent concentration of effluent i, e effluent with-

out dilution causes decrease in vegetative growth in comparison to control condition due to presence of sulphate and phosphate in the effluent. Vegetative growth increases with the increase in concentration of effluent upto 80 percent.

## REFERENCES

- Aggarwal, R.K., Sharma, B.M. and Kumar, P. (1989). Potential use of fly ash for improving physico- chemical properties of sandy soils. *Int. Symposium Managing Sandy Soils. Part – I*: 133-138.
- Baruah, B.K. and Dass, M. (1998). Study the impacts of paper mill effluent on germination behavior and seedling growth of crop plant *Oryza sativa L.* *Pollution Resources*,17(1): 65-68.
- Dutta, S.K., Boissya, C.L. (1997). Effect of paper mill effluent on germinations of rice seed (*Oryza sativa L.* var. Masuri) and growth behaviour of its seedlings. *J Indl Polln Contl*, 13(1), 41-47.
- Prasanna Kumar, P.G., Pandit, B.R., Mahesh Kumar, R. (1997). Effect of dairy effluent on seed germination, seedling growth and pigment content of green gram (*Phaseolus aureus L.*) and black gram (*Phaseolus mango L.*). *Adv Plant Sci*, 10(1), 129-136.
- Ramana, S., Biswas, A.K., Kundu, S., Yadav, R.B.R. (2002) Effect of distillery effluent on seed germination in some vegetable crops. *Bioresource Technology*, 82, 273-275.
- S. Arora, A.K. Chopra *et. al.*, (2005).



Physicochemical and bacteriological characteristics of Aachal Dairy milk effluent and its effects on seed germination of some agricultural crops. *Nature, Environment and Pollution Technology*. 4(3): 441-444.

7. Singh, Anoop, Agrawal, S.B., Rai, J.P.N., Singh, Pratibha (2002). Assessment of the pulp and paper mill effluent on growth, yield and nutrient quality of wheat (*Triticum aestivum* L.). *J Environ Bio*, 28(3), 283-288.
8. Singh, G. and Gupta, S.K. (1989). Potentiality of fly ash in augmenting the physico- chemical properties of sandy soils for improved crop- production. *Int. Symposium Managing Sandy Soils Part I*. 142-145.
9. Singh, Lamabam P., Siddiqui, Zaki A. (2003). Effects of fly ash and *Helminthosporium oryzae* on growth and yield of three cultivars of rice. *Biore-source Techno*, 86(1), 73-78.
10. Yadav, J.P. and Minakshi (2006). Effect of sugar mill and milk plant effluent on the seed germination and early seedling growth of agricultural crops. *Pollution Resources* 25(4): 701-705.
11. Young, J.C., Boumann, E.R. and Danyal, V. (1975). Packed bed reactors for secondary effluent BOD and ammonia removal. *J. Wat. Pollut. Control Fed.*, 47, 46.







# STUDY THE EFFECT OF DAIRY EFFLUENTS ON SEED GERMINATION OF HIGH NUTRIENT QUALITY PEARL MILLET-*Pennisetum typhoides*(L.)

**Goutam Bhakta**

Rukminipur K.L Vidyapith (High),  
Egra, Purba Medinipur, West Bengal

Email : [ani.ezee@gmail.com](mailto:ani.ezee@gmail.com)

(Date of Receipt :26-08-2014 ;

Date of Acceptance for Publication : 23-09-2014)

## Abstract

Today environmental degradation has become a global phenomenon due to industrialization and urbanisation. Most of the effluents contain varied groups of chemical compound including nutrients. This nutrient helps in fertilization of soil and would also increase productivity of the land. The present study deals with the effect of dairy effluent on seed germination of the crop high nutrient pearl millet *Pennisetum typhoides*(L) : The main objective of this study is the effect of dairy effluents on seed germination. Percentage of pearl millet in various concentrations of effluent (20%, 40%, 60%, 80%, 100% and control in laboratory in Petridis. The germination percentage of pearl millet decreased as the concentration of effluent increased.

**Key words** :- Pearl Millet, Dairy Effluent, Germination.

**Pages: 4**

**References: 9**

## INTRODUCTION

Environmental degradation has now become a global problem and maintaining ecosystem health is a serious issue by environmentalists. Due to lack of effluent treatment facilities and proper disposal system of waste, water bodies are getting polluted day by day and causing adverse effect on soil, water, agriculture, flora and fauna due to presence of toxic and persistent chemicals. So it become essential either to find suitable ways for the safe disposals of these wastes or to suggest their novel use by their by-product. Finding a profitable use for this waste could

further benefit the economics of industry. Among these industries food processing industrial effluents relapsed from the dairy industry Milk Plant, Sirsa. Haryana is rich in phosphate, calcium and magnesium etc. and has good potential in utilization of released effluents as a source of nutrients for the crop plants like *Pennisetum typhoides* (Pearl millet). Pearl millet is a high nutrient quality having high amount of protein fat, a basic staple for households in the poorest countries and among poorest people. For this, studies related of seed germination were carried out to check the viability of seeds. In my

present work I have investigated the germination percentage of seed pearl millet seed along with radicle and plumule lengths and also the fresh weight and dry weight of germinated seedlings.

## MATERIAL AND METHODS


To carry out the study for physico chemical analysis of soil collected from Balsamand area, Hisar air dried soil samples passing through 2 mm sieve were used, pH, electrical conductivity, calcium carbonate, organic carbon, nitrogen, phosphorous, potassium are done and for physico chemical analysis of dairy effluents pH, electrical conductivity, total dissolved solids, calcium, Magnesium, sulphate, phosphate, chloride, BOD, COD are estimated. To carry out the effect of dairy effluent seed germination experiments first to check the viability of seeds. For this twenty seeds of pearl millet were soaked in water taken in a beaker for 30 minutes. After that these seeds were placed on double layered water soaked filter paper placed in petri plates. The covered petriplates were kept in BOD at 25+ 1°C for 3 days. After 3 days, first observation was done for their viability; secondly to study the effect the effect of various concentrations of effluents (20%, 40%, 60%, 80%, 100% and control) of milk plant on seed germination percentage, first all seeds of pearl millet sterilized in  $HgCl_2$  for 2 minutes. Eighteen petriplates (for six setups in multiple of three) were used. At one time ten seeds were taken on double folds of whattman no. 1 filter paper for every petriplate and were supplied with different dose of effluents in controlled condition at 25+1°C in seed germination for the experimental period

1–8 days Reading were taken after. 3<sup>rd</sup> day, 5<sup>th</sup> day and 8<sup>th</sup> day. On 8<sup>th</sup> day radical and plumule lengths were noted and radical / plumule ratio was derived; dry and fresh wt of germinated seedlings were estimated.

## RESULTS AND DISCUSSION

### Bar 1.

### RESULTS OF SEED GERMINATION EXPERIMENT



The germination percentage of pearl millet decreased as the concentration of effluent increased (Fig.: 4.3.1). The maximum germination percentage i.e. 100 per cent was observed under control and minimum i.e. 63.3 per cent was observed under 100 per cent concentration of effluent on 8<sup>th</sup> day of germination. The percentage of seed germination increased up to 60 per cent concentration of effluent. The percentage of seed germination at 20 per cent concentration of effluent was found to be 80 per cent and at 60 per cent concentration it was found to be 90 per cent. The similar observations were recorded by (Gautam *et. al.*, 1992; Arora *et. al.*, 2005; Ajmal *et. al.*, 1984). The viability of the seeds was recorded to be 100 percent.

Table 1.

Effect of different concentrations of dairy/milk plant effluent on the plumule length and radical length(cm) of germinated seedlings of pearl millet after 8 days.

Concentration	Length of Plumule (cm)	Length of Radicle (cm)	Plumule : Radicle Ratio
Control	5.2	16.2	0.32
20%	5.3	16.4	0.32
40%	6	17.3	0.35
60%	6.5	18.3	0.35
80%	3.4	12.5	0.27
100%	1.3	4.9	0.26

Plumule radicle ratio was recorded maximum i.e. 0.35 at 40 percent & 60 percent conc. of effluent and minimum i.e. .26 at 100 percent effluent.

Table 2:

Effect of different concentrations of dairy/milk plant effluent on the fresh weight and dry weight(gm) of germinated seedlings of pearl millet after 8 days.

Concentrations	Fresh wt. of germinated seedlings (gm)	Dry wt. of germinated seedlings (gm)
Control	1.291	0.109
20%	1.319	0.119
40%	1.411	0.127
60%	1.450	0.138
80%	0.954	0.089
100%	0.540	0.047

The fresh wt. and dry wt. of germinated seedling of pearl millet were observed to be minimum i.e. 0.540 gm and .047 gm respectively at 100 percent concentration; and to be maximum i.e. 1.450 gm and 0.138 gm respectively at 60 percent conc. of effluent.

## CONCLUSION

It is concluded that germination percentage of pearl millet decreased as the concentration of effluent increased i.e.

i) 100 percent germination was observed under control and minimum under 100 percent concentration of effluent.

ii) Plumule radicle ratio was recorded maximum at 40 percent and 60 percent and minimum at 100 percent concentration of effluents.

iii) Fresh wt. and dry wt. of germinated seedling of pearl millet also increased as the conc. of effluent increased upto 60 percent and decreased upto 100 percent conc. of effluent.

## REFERENCE

1. Aggarwal, R.K., Sharma, B.M. and Kumar, P. (1989). Potential use of fly ash for improving physico- chemical properties of sandy soils. *Int. Symposium Managing Sandy Soils. Part – I*: 133-138.
2. Baruah, B.K. and Dass, M. (1998). Study the impacts of paper mill effluent on germination behavior and seedling growth of crop plant *Oryza sativa* L. *Pollution Resources*,17(1): 65-68.
3. Dutta, S.K., Boissya, C.L. (1997). Effect of paper mill effluent on germinations of rice seed (*Oryza sativa* L. var. Masuri) and growth behaviour of its seedlings. *J Indl Polln Contl*, 13(1), 41-47.
4. Prasanna Kumar, P.G., Pandit, B.R., Mahesh Kumar, R. (1997). Effect of dairy effluent on seed germination, seedling growth and pigment content of green gram (*Phaseolus aureus* L.) and black gram (*Phaseolus mango* L.). *Adv Plant Sci*, 10(1), 129-136.



5. Ramana, S., Biswas, A.K., Kundu, S., Yadav, R.B.R. **(2002)** Effect of distillery effluent on seed germination in some vegetable crops. *Bioresource Technology*, 82, 273-275.
6. S. Arora, A.K. Chopra *et. al.*, **(2005)**. Physicochemical and bacteriological characteristics of Aachal Dairy milk effluent and its effects on seed germination of some agricultural crops. *Nature, Environment and Pollution Technology*. 4(3): 441-444.
7. Singh, Anoop, Agrawal, S.B., Rai, J.P.N., Singh, Pratibha **(2002)**. Assessment of the pulp and paper mill effluent on growth, yield and nutrient quality of wheat (*Triticum aestivum* L.). *J Environ Bio*, 28(3), 283-288.
8. Yadav, J.P. and Minakshi **(2006)**. Effect of sugar mill and milk plant effluent on the seed germination and early seedling growth of agricultural crops. *Pollution Resources* 25(4): 701-705.
9. Young, J.C., Boumann, E.R. and Dan-nyal, V. **(1975)**. Packed bed reactors for secondary effluent BOD and ammonia removal. *J. Wat. Pollut. Control Fed.*, 47, 46.







Bimonthly Newsletter for Scientific Community

**Academic & RESEARCH Publications**  
ARP Give your JDCoA an Expression

**BEST**  
(BULLETIN ON EMERGING SCIENCE AND TECHNOLOGY)  
PUBLISHED BY Academic and Research Publications  
22, Gaur Galaxy, plot no 5, sector 5,  
Vaishali Ghaziabad UP 201010

The main aims and objectives of our publishing house are to Publish newsletter, magazine, journals, books etc and to organize seminar, symposium, conferences etc for the dissemination of knowledge from local to global.

SCIENTIFIC COMMUNITY NEWS LETTER


Vol. 1, Issue1, Dec 2014  
FOR INTERESTED SCIENTISTS ONLY

**NEWSletter**

RECENT TRENDS IN DEVELOPMENT OF SCIENCE AND TECHNOLOGY FOR HUMAN BENEFITS

## BEST (BULLETIN ON EMERGING SCIENCE AND TECHNOLOGY)

### European Space Agency's Goc satellite falls to Earth



Goc needed its engine power to stay in orbit



*Prof. T.R.C Sinha a true lover of science and always concerned about the environment. It was his dream to publish scientific journals and he started this project with the same enthusiasm as he had done for every project he undertook. We are guided by the vision of Prof. Sinha and endeavour to make his dream of promoting science and help young scientists, publish their articles and encourage them in their research.*

*Manisha Verma*

 Submit your manuscripts

@ manisha\_npp  @ yahoo.com



manishanpp

Academic and Research Publications







**International Journal  
Of  
Agricultural Science & Technology**  
(Bi-Annual Journal)  
2014

**SUBSCRIPTION ORDER FORM**

Name: \_\_\_\_\_

Address: \_\_\_\_\_

Pin: \_\_\_\_\_

Tel: \_\_\_\_\_ Fax: \_\_\_\_\_

E-mail: \_\_\_\_\_

**MODE OF PAYMENT**

**Payment should be made online. The amount is to be credit in the Bank The details of the bank are as follows:-**

Name of the Account holder : **Academic And Research Publications**

Name of the bank : **PUNJAB NATIONAL BANK, VAISHALI BRANCH, GHAZIABAD,  
IFSC: PUNB0405300**

Current Account No. **4053002100011106**

New ☐

Renewal ☐

Signature .....Date .....

Published by

**Academic And Research Publications**

Office: 22, Gaur Galaxy, Plot No 5, Sec-5, Vaishali , Ghaziabad (U.P.) - 201010 (INDIA)



# National Journal of Comparative Law

## Brochure

[www.manishanpp.com](http://www.manishanpp.com)





## SUBSCRIPTION RATES

### INDIVIDUAL COPY for CO-AUTHORS Only

<b>Inland</b>	<b>Foreign</b>
<b>Rs. 500/-</b>	<b>US\$ 40</b>
<b>ANNUAL SUBSCRIPTION</b>	
<b>(2 Issues Per Year)</b>	

<b>Individuals</b>	<b>Foreign</b>
<b>Rs. 800/-</b>	<b>US\$ 80</b>

<b>Institutions</b>	<b>Foreign</b>
<b>Rs.1500/-</b>	<b>US\$ 100</b>

## ADVERTISEMENT RATES



### Full Page

Colour	:	Rs. 35,000.00
Black/White	:	Rs. 25,000.00

### Half Page

Colour	:	Rs. 25,000.00
Black/White	:	Rs. 15,000.00

**(Four weeks to be allowed after submission of advertising material)**



**Submit your manuscripts**

@ [manisha\\_npp@yahoo.com](mailto:manisha_npp@yahoo.com), [arp@manishanpp.com](mailto:arp@manishanpp.com)

[www.manishanpp.com](http://www.manishanpp.com)

Copyrights@www.manishanpp.com 2015- All rights reserved

## Contact Information

For Subscription orders may be contact directly to publisher (address given below) using Journal Subscription Form or contact your preferred subscription agent. All orders should be accompanied by payment. Please make checks payable to ("Academic And Research Publications"). Mail order with payment to:

Subscription Manager :

### Academic And Research Publications

**Office:** 22, Gaur Galaxy,  
Plot No 5, Sec-5, Vaishali ,  
Ghaziabad (U.P.) - 201010 (INDIA)

Email: [manisha\\_npp@yahoo.com](mailto:manisha_npp@yahoo.com)

Phone No.: 0120-4124773

For more information or assistance to place an order, please feel welcome to contact us again at

[manisha\\_npp@yahoo.com](mailto:manisha_npp@yahoo.com)

## Mode of Payment

Subscriptions are payable in advance and all rates include postage and taxes. Subscribers are requested to send payment with their order whenever possible. Issues will only be sent on receipt of payment. Subscriptions are entered on an annual basis – January to December – and paid in Rs or US Dollars. For payments in Euros and Sterling, please calculate at the prevailing exchange rate.

Payment can be made by Bank (Wire) Transfer, Cheque, International Money Orders. All payments should be made to "Academic And Research Publications".

Join us on facebook



ARP



## JOURNAL SUBSCRIPTION ORDER FORM FOR 2015

Name: \_\_\_\_\_ Dept. \_\_\_\_\_  
Address: \_\_\_\_\_  
Tel: \_\_\_\_\_ Mob. \_\_\_\_\_  
E-mail: \_\_\_\_\_ Pin: \_\_\_\_\_

- ☒ Check appropriate boxes (Annual Subscription Price in 1000 Only for *Institution*)
- ☐ National Journal of Comparative Law(NJCL) ISSN : 2393-9338
- ☐ International Journal of App. Environmental Science & Technology : 2321-8223
- ☐ International Journal of Agricultural Science And Technology : ISSN 2319-880X
- ☐ International Journal of Aquaticscience And Technology : ISSN 2320-6772
- ☐ International Journal of Nanoscience And Technology : ISSN 2319-8796
- ☐ International Journal of Geoscience And Technology : ISSN 2321-2144

### MODE OF PAYMENT

Payment should be made online. The amount is to be credit in the Bank The details of the bank are as follows:-

Name of the Account holder : Academic And Research Publications

Name of the bank : PUNJAB NATIONAL BANK, VAISHALI BRANCH,  
GHAZIABAD,

IFSC: PUNB0405300

Current Account No. 4053002100011106

New

☐

Renewal

☐

SUBSCRIBER TYPE: (Check one) Library ☐ /Institution ☐ / Personal ☐

Signature

..... Date .....

Published by  
**JPMS Society**

Office: 22, Gaur Galaxy, Plot No 5, Sec-5, Vaishali , Ghaziabad (U.P.) - 201010 (INDIA)

www.manishanpp.com

Join us on  ARP

To order by telephone, please call us at +0120-4124773

Sample Issue: To access free online sample issue of our journals, please send e-mail to [manisha\\_npp@yahoo.com](mailto:manisha_npp@yahoo.com)





*Dear Reader,*

*This journal is a collaborative effort which aims to increase awareness and understanding of scientific research, This is a peer reviewed biannual journal which gives a platform to fresh ideas and past experience about developements in India and outside. The journal has Open Access to its content and makes knowledge available to the public in general and re-search scholars in specific. Through this journal we welcome greater global exchange of knowledge, ideas among intellectu-als, researchers and scholars. The journal offers [an open forum for discussion](#). I will like to appreciate the efforts of all authors, co-authors, reviewers and member of Editorial Board who have contributed immensely towards this journal.*

*Manisha Verma*

**Copyright: Submission of a manuscript implies: that the work described has not been published before (except in the form of an abstract or as part of a published lecture, or thesis) that it is not under consideration for publication elsewhere; that if and when the manuscript is accepted for publication, the authors agree to automatic transfer of the copyright to the publisher.**

---

# NOTES



## NOTES



## Guidelines to Authors

### International Journal of Agricultural Science & Technology

This journal is published by the **Academic And Research Publications New Delhi**, twice in a year. The emphasis is to involve a large community of scientists specially JUNIOR SCIENTISTS and scholars from India and abroad in developing a framework of discussion and debate on conservation and sustainable development. Research articles, Review articles, Book reviews, Interviews, Short communication, Letters to the Editor, Case reports and News items related to the subject are accepted for publication. The PUBLISHER is aiming to publish your article as rapidly as possible provide a rapid publishing platform, each and every point is vital to reduce the editorial work flow. Authors are requested to extend their maximum cooperation for speedy actions to be taken at this end. Hence they are requested to check the guidelines thoroughly before submitting their article(s) and check that all the required informations are mentioned and strictly according to the guide lines.

### Benefit to Authors

Following benefits shall be available to the authors, if your article is published in the journals of this PUBLISHER. Authors can use their published articles in any way they like for the dissipation of knowledge free of charge. However it cannot be used for any commercial use without prior permission. Authors can have access on all articles published by this Publisher. Wide publicity and reach by getting indexed in many free indexing services.

### Submission of Article(s)

All manuscripts are to be submitted in English, typed double-spaced throughout the text. It is requested that manuscripts be sent by e-mail only. Authors will be requested to substantiate the need if it exceeds the maximum number of pages.

All manuscript should be submitted to the Publication Editor at [manisha\\_npp@yahoo.com](mailto:manisha_npp@yahoo.com) directly.

### The Research Article should be in the following order

#### In the 1st Part:

1) Title 2) Name of the organisation where work was carried out 3) Name of the Author with whom correspondence is to be made along with his/her Mob.No. 4) Date of Submission 5) No. of Tables 6) No. of Photographs 7) Numbers of Graphs 8) No. of References 9) Name and addresses of at least Three referees along with their Email I/Ds and Mob Nos.

#### In the 2nd Part :

1) Title of The Article 2) Name and addresses of all the Author(s) along with their Email I/Ds and Mob.Nos. 3) Abstract 4) Introduction 5) Materials and methods 6) Results 7) Discussions (if any) 8) Acknowledgements 10) References (Under References heading, Name of the Journals should be in bold letters) and (11) Name of The journal.

### Tables

to be included should have a heading, giving the substance, and should be typed double-spaced on separate sheets. They should also be numbered in serial order. Figures either drawn manually or by computer should be in black ink and the lettering on them should be large enough to stand reduction. Photographs in colour should have sharp contrast. Legends for figures and plates should be typed in numerical order on separate sheets, one for figures and one for plates.

### References

The literature cited should list the author's name, year of publication, title of the paper, and the Journal titles (bold letters) which should be cited in full (no abbreviation) with volume number and page numbers, as indicated below:

**A. For articles in a Journal:**-Walsh, J.E. (2008) Climate of the Arctic Marine Environment. Ecological Applications. 18. pp. 3-22.

**B. For Books:**-Ward, D.R. (2002) Water Wars: drought, floods, folly and politics of thirst: River head Books. New York. p. 12.

**C. Chapter in a book:**-Andrews, T.J., Cough, B.F. and Muller, G.J. (1984). Photosynthetic gas exchange properties and carbon isotope ratios of some mangroves in North Queensland. In: H.J. Teas (Ed.), Physiology and Management of Mangroves. W. Junk. The Hague. pp. 15-23.

**From website:**-National Oceans and Atmospheric Administration (NOAA). 1995. Regional Perspectives: Indian Ocean. [www.ncdc.noaa.gov/paleo/outreach/coral/sor/sor\\_indian.html](http://www.ncdc.noaa.gov/paleo/outreach/coral/sor/sor_indian.html), accessed on July 13, 2008.

While giving reference of more than two authors in the text, after the name of the first author, *et.al.*, should be used, followed by the year of publication.

### PRINTING CHARGE

The articles are not printed in gratis. A nominal charge of Rs 500=00 per printed page is charged and is printed in B/W colour. For any coloured page, is separately charged. Charges for the same is quoted on request. Five Reprints and a copy of Journal is provided to the Corresponding author. Extra copies of the Journal for coauthor only are supplied on charge @ Rs 400-00 per copy and shall be sent along with the copy of the Correspondent Author.

### REQUEST

Authors are requested to keep a copy of the Mss. till it is published in the Journal.

### REMITTANCE

All remittances are to be on line credited in the account of **Academic And Research Publications** in the **PUNJAB NATIONAL BANK, VAISHALI BRANCH, GHAZIABAD, IFSC: PUNB0405300**

**Current Account No. 4053002100011106** and inform the at [manisha\\_npp@yahoo.com](mailto:manisha_npp@yahoo.com) accordingly.

### CONTACT US

For quick reply, please note the address and contact them directly by Post or email:-a) For publication of your article, Acceptance letter, Invoice, sending of Cheques/Drafts, for sending Review Reports, Status Report about your article, and all other queries related to your articles, should be sent directly to the Editor-in-Chief, whose address is as follows:

**Prof. Manik Sinha-THE EDITOR-IN-CHIEF, H.Office: EC 41, Maya Enclave, New Delhi -110064**

Email: [manik.sinha2@gmail.com](mailto:manik.sinha2@gmail.com), [manisha\\_npp@yahoo.com](mailto:manisha_npp@yahoo.com). [www.manishanpp.com](http://www.manishanpp.com).




## Contents

### *Reviewed Articles*

1. Allelopathic effect of *Eucalyptus tereticornis* clone3 on germination and growth of *Vigna mungo* L 1  
V. D. Tripathi, A. Venkatesh and S. K. Dhyani
2. Use of Medicinal plants for treatment of cancer-A review 11  
Malika Pal
3. Effect of Dairy Effluents on Early Seedling Growth of *Pennisetum Typhoides* (L) 17  
Goutam Bhakta
4. Study The Effect of Dairy Effluents on Seed Germination of High Nutrient Quality Pearl Millet-*Pennisetum Typhoides*(L.) 22  
Goutam Bhakta

An International Refereed Journal

**Academic & RESEARCH Publications**  
ARP. Give your JDEA an Expression  
[www.manishanpp.com](http://www.manishanpp.com)



**ACADEMIC AND RESEARCH PUBLICATIONS**

H.Office: EC 41, Maya Enclave,  
New Delhi -110064

Email : [manik.sinha2@gmail.com](mailto:manik.sinha2@gmail.com)  
[manisha\\_npp@yahoo.com](mailto:manisha_npp@yahoo.com)

Join us on facebook  ARP